

INDONESIAN JOURNAL OF SOCIAL AND ENVIRONMENTAL ISSUES (IJSEI)

Journal Homepage: https://ojs.literacyinstitute.org/index.php/ijsei ISSN: 2722-1369 (Online)

Research Article

Volume 2	Issue 3	December (2021)	DOI: 10.47540/ijsei.v2i3.281	Page: 204 – 212
		× /	5	U

Biomass and Carbon Stock in the Sal (Shorea robusta) Forest of Dang District Nepal

Supuspa Regmi¹, Krishna Prasad Dahal², Garima Sharma³, Siddhartha Regmi⁴, Mahamad Sayab Miya⁵

¹⁻⁵Institute of Forestry, Tribhuvan University, Nepal

Corresponding Author: Mahamad Sayab Miya; Email: sayabmiya13@gmail.com

ARTICLEINFO

ABSTRACT

Keywords: Biomass; Carbon Stock; The study was conducted to find the net carbon stock in the Sal (Shorea robusta) Growth Modeling; Shorea robusta; forest in Bagdaila Chisapani community forest of Dang district, Nepal. The Tropical Forest. inventory was done by a stratified sampling technique with 0.5% sampling intensity taking into account the woody plants with \geq 5cm DBH. A total of 49 sample plots Received : 25 June 2021 of radius 8.94m and 5.64 were established to measure tree biomass and sapling Revised : 11 December 2021 biomass respectively in the forest. The mean above-ground carbon (AGC) was : 14 December 2021 Accepted 160.4 t ha⁻¹ and the mean below-ground carbon (BGC) was 24.1t ha⁻¹. The mean total carbon stock in the study area was estimated to be 99.02 t ha⁻¹; of which maximum carbon stock of 143.51 t ha⁻¹ was found in block number 5. Sal was the major tree species in the CF. The biomass and carbon content in this forest is found quite low as compared to other studies in the Sal forest and other tropical forests which were due to the presence of tree stands of less diameter and height. This study would be helpful in the long-term management of forests, planning, and research purposes. The data from this study could also be taken as a reference document for the participation of community forests in carbon accounting under the REDD+ scheme in Nepal.

INTRODUCTION

Forests store huge amounts of the global carbon (terrestrial above-ground carbon-80% and below ground carbon-40%) and play a significant function in global carbon balance and cycle (Dixon et al., 1994; Kirschbaum, 2000; Eggleston et al., 2006; Candell and Raupach, 2008; Pan et al., 2011). In the forests, carbon is typically reserved in tree trunks, leaves, foliage, and roots referred to as biomass. And they act as a natural brake for climate change and related issues (Gibbs et al., 2007; Fahey et al., 2010). Carbon is generally accumulated in biomass during the photosynthesis process (Alexandrov, 2007; Suryawanshi et al., 2014). All parts of a plant contain carbon, but the proportion in each part varies extremely. The carbon stock (CS) and biomass in the forests usually differ with species, type, age, canopy cover, stand structure, and altitude (Pandey et al., 2014; Karki et al., 2016; Dar et al., 2017). After the death of plants, biomass reaches the soil as soil carbon. When biomass is destroyed, the carbon is re-discharged to the atmosphere (Banik et al., 2018). Forest biomass provides opportunities for ecological visioning, sustainable forest management, enhancement of ecosystem functioning and services (Pan et al., 2013; Panzou et al., 2018), and help in climate change mitigation (Houghton, 2005; Eriksson and Berg, 2007).

The Kyoto Protocol was formulated in 1997 stating the principle that the soil and biomass can sequestrate CO_2 from the air; which is the real solution for climate change (Harishma et al., 2020). Stopping deforestation alone will lead to a reduction of about 18% of atmospheric CO_2 emissions (IPCC, 2007). Emissions reduction can be achieved through proper management of forests with focused objectives of REDD+ programs (Skutsch and Laake, 2008). According to FAO (2011), about 7% of the forests in the world are managed by communities under community forest management programs. These community-managed forests have enough potential to store carbon (Tripathi et al., 2017).

Tropical forests of the world cover 7-10% of the global land (Raha et al., 2020) but store more than 40% of the terrestrial carbon in the form of biomass (Phillips, 1988). These forests contribute about 34% of the primary productivity in terrestrial lands (Lewis et al., 2013). Tropical forests of Nepal are dominated by *Shorea robusta* (Sal) (Jackson, 1994). It is distributed from Terai to 1500m elevation from sea level (Gautam and Devoe, 2006). Sal covers more than half of the tropical forests in the Terai region (Webb and Sah, 2003). The *S. robusta* forests of Nepal are among the important carbon-storing tropical forests (Shrestha, 2008).

Above-ground biomass (AGB) provides crucial information on several global challenges (Brown et al., 1999). Carbon stock is estimated based on AGB (Ketterings et al., 2001; Cao et al, 2001; Mokany et al., 2006). Carbon storage estimation help to meet information related to greenhouse gas (GHG) reduction (Adme et al., 2020). It is also needed to implement climate change-related strategies and programs (eg. REDD+) (Saatchi et al., 2011; Avitabile et al., 2016). Kyoto Protocol, Paris Agreement, and European Union (Regulation 2018/841) have also appealed to estimate, document, and report carbon, carbon emissions, and extent of deforestation (UNFCCC, 2008; UNFCCC, 2015; Forsell et al., 2018). But, in developing countries, the biomass and carbon stocks in tropical forests are feebly estimated (Behera et al., 2017) and limited due to the lack of atmospheric observations (Kaul et al., 2010; Reichstein et al., 2013; Mitchard et al., 2014).

Also in Nepal, very little researches have been conducted relating to biomass and carbon stock in tropical forests. However, several studies are conducted in various regions of the country (Upadhyay et al., 2005; Nepal, 2006; Baskota et al., 2007; Baral et al., 2009; Subedi et al. 2010; Bhusal, 2011; Pandey and Bhusal, 2016). This study was performed in one of the tropical forests of Dang district, Nepal with the aims of estimating the AGC and BGC and intimately finding out the carbon stocks/content based on the biomass.

MATERIALS AND METHODS Study area

The study was carried out in Bagdaila Chisapani community forest (BCCF) of Dang district (82[°] 2' to 82[°] 54'E and 27[°]37' to 28[°] 36'N) (Figure 1). The district lies in the inner Terai region of Nepal and borders India in the south. It covers an area of 290,251 ha with an altitude range of 213m to 2058 m above sea level. Geographically, the district is divided into two parts i.e. Plain valley (Bhabar) and Siwalik Hills. It is bounded by the outer Siwalik Hills in the south and the Mahabharat hills in the north. Three types of climatic zones are observed in the district i.e. lower tropical (<300m, occupy 18.1%), upper tropical (300-1000m, occupy 69.9%), and subtropical zone (1000-2000m, occupy 12%). The annual precipitation of the district varies between 1584 mm and 2287 mm. The soil is dominated by clay and sandy loam (Pandey et al, 2019). Forests cover the maximum area of the district (over 50%) and the remaining land use are agriculture, water body, and others (Ghimire, 2017). The total population of the district is 552,583 (CBS, 2012). BCF lies in Babai rural municipality. The forest has a total area of 254.6 ha dominated by Shorea robusta. Scientific forest management program has been applied in the community forest since 2015; hence the total forest area was assigned into eight blocks.



Figure 1: Study area map showing Bagdaila Chisapani CF

Data collection

The primary data was obtained through stratified sampling. The already divided forest blocks during scientific forest management plan preparation were taken as the strata. Total 49 circular plots (0.5% sampling intensity) were prepared in different blocks of the forest. Aboveground biomass for (DBH \geq 5 cm) was measured in the circular plots (r=8.92m). While, above-ground sapling biomass (DBH= 1-5cm) was measured inside nested plots (r=5.64m) (ANSAB, 2010). The total height (H) and diameter at breast height (DBH) of entire trees having DBH≥ 5 cm were measured carefully inside the plots. Secondary data was gathered from several documents obtained and accessed through Google Scholar and Research Gate.

Data analysis

Microsoft Excel was used for the database, simple modeling, and descriptive statistics like percentage, mean and median. The variability in biomass was measured and expressed in the form of the SE (standard error of the mean).

Above-ground tree biomass (AGTB)

It was estimated with the simplified standard regression model where DBH (D), Height (H), and wood density/specific gravity (ρ) were considered for calculation (Chave et al., 2005; Vachnadze et al., 2018; Komolafe et al., 2020).

$$4GTB = 0.0509 * \rho D^2 H$$

(Here, AGTB is in kg, ρ is in g cm⁻³, D in cm, and H in m). The obtained AGTB value for all individuals of a sapling plot was summed up; then divided by 250 m² (sampling plot area). And it was multiplied by 10 to convert into t ha⁻¹.

(1)

Above ground sapling biomass (AGSB)

It was estimated with the formula given by (Tamrakar, 2000); which is the logarithmic transformation of the allometric formula.

$$Log (AGSB) = a + b log (D)$$
(2)

[AGSB is in kg, a = intercept of algometric relationship (unitless), b = slope algometric relationship (unitless), D is in cm]. (a= -2.4554, b= 1.9026, and R^2 = 98.3).

Below-ground biomass (BGB)

 $BGB = 0.15 \times above$ -ground biomass (AGB) (3)

Net carbon stock (CS)

It was estimated with the stock method where the total carbon stock/content is supposed to be 47% of the total dry biomass (DB) (Eggleston et al., 2006).

CS = Total DB of the tree x 47% (4)

RESULTS AND DISCUSSION Forest Biomass

The BCF was divided into a total of eight blocks. The average total biomass of the BCCF was Table 1. Biomass in different blocks 184.4 t ha⁻¹. There was an average of 263 stems ha⁻¹ with a mean DBH of 17.1 cm. The highest number of stems per ha was found at block number 3 followed by block number 7. There was an average of 1421 saplings ha⁻¹ with a mean DBH of 4.2 cm. Block number 5 has the highest tree biomass (305.3 t ha⁻¹) followed by block numbers 4, 3, 7, 1, 8, 2, and 6. The result for the total AGB and BGB also follows the same trend in different blocks (Table 1).

	Tree characteristics				Sapling characteristics			AGB		BGB		
Blocks	No of Stems Per ha	Mean DBH (cm)	Mean Height (m)	Tree Biomass (t/ha)	SE	No of Sapling Per ha	Mean dbh (cm)	Sapling Biomass (t/ha)	Total AGB (t/ha)	SE	Below ground biomass (t/ha)	Total Biomass
1	286	17.3	11.5	154.2	12.3	1254.0	4.9	0.95	155.1	12.3	23.3	178.4
2	285.0	14.8	10.5	103.4	14.5	2123.9	4.6	0.98	104.3	14.5	15.6	119.9
3	305.0	18.5	11.6	177.9	15.2	1256.0	4.8	0.82	178.7	15.2	26.8	205.6
4	215.0	19.2	11.5	189.9	9.6	944.0	3.8	0.50	190.4	9.6	28.6	219.0
5	236.0	20.3	14.3	264.7	7.2	1434.0	4.1	0.83	265.5	7.2	39.8	305.3
6	252.0	13.3	11.5	91.7	13.8	1345.0	3.1	0.49	92.1	13.8	13.8	105.9
7	304.0	17.6	12.2	169.3	10.5	1523.5	4.3	0.83	170.1	10.5	25.5	195.6
8	222.0	15.8	11.3	125.9	13.0	1492.0	4.2	0.81	126.6	13.0	19.0	145.6
Average	263.6	17.1	11.8	182.4	12.0	1421.0	4.2	0.7	160.4	12.0	24.1	184.4

Total carbon stock

The total CS in the BCCF was 99.02 t ha⁻¹. The highest CS was in block number 5 (143.51 t ha⁻¹) Table 2. Carbon content in a different block

followed by block numbers 4, 3, 7, 1, 8, 2, and 6. The area of the BCCF is 254.6 ha. Thus, the total CS is estimated to be 25,210.49 tones (Table 2).

Blocks	Ab	ove Ground	Carbon	Below Ground	Total Carbon	
	Tree (t ha ⁻¹)	Sapling (t ha ⁻¹)	Total (t ha ⁻¹)	SE	Carbon (t ha ⁻¹)	(t ha ⁻¹)
1	72.46	0.44	72.90	5.80	10.93	83.83
2	48.61	0.40	49.01	6.83	7.35	56.37
3	83.61	0.39	84.01	7.16	12.60	96.61
4	89.25	0.24	89.49	4.53	13.42	102.91
5	124.43	0.37	124.79	3.38	18.72	143.51
6	43.08	0.20	43.29	6.47	6.49	49.78
7	79.56	0.39	79.94	4.91	11.99	91.93
8	59.16	0.35	59.51	6.10	8.93	68.44
Average	85.74	0.35	86.08	5.65	12.91	99.02

The CS and biomass in the forests differ with species, type, age, canopy cover, stand structure, and altitude (Pandey et al., 2014; Karki et al., 2016; Dar et al., 2017). In the present study, the total CS is varied from block to block. A study carried out by Gautam and Mandal (2016) in the Sunsari

district found that the plant biomass for the undisturbed forest was greater (960.4 t ha^{-1} or 452.06 t C ha^{-1}) than the disturbed forest (449.1 t ha^{-1} or 211.33 t C ha^{-1}); where the stand density for the disturbed forest was 234 stems ha^{-1} . (Ngo et al., 2013) have found the total carbon content higher in

the primary forest (337 t C ha⁻¹) than in the secondary forest (274 t C ha⁻¹). Mandal et al., 2013) has found CS higher in Gadhanta-Bardibas CFM $(274.66 \text{ t ha}^{-1})$ than in Banke-Maraha CFM (197.10) t ha^{-1} ; both of the forests have *Shorea robust* as the dominants species in the Mahottari district of Nepal. Bohora et al. (2021) has estimated the total biomass of 1381.30 t ha⁻¹ and CS of 649.21 t ha⁻¹ in Shorea robusta and Schima wallichi forest of Makwanpur district. In the research conducted by Ghimire (2017) in the Danphe community forest of Dang district, the average CS was 62.34 t ha⁻¹ in 2013 whereas; it was 64.86 t ha⁻¹ in the year 2014. The CS is greater in this study as compared to Ghimire (2017). In the Kayerkhola watershed of Chitwan district (dominated by the Shorea robusta forest), the sparse area had 89.2 t C ha⁻¹ while the dense area had 129.0 t C ha⁻¹ (Pandey et al., 2014). The estimated total biomass and CS are quite higher than in our study. Our study area had an immature forest having tree species with small DBH and height which is 17.1 and 11.8 respectively on average. This might be the cause for lower biomass and CS in BCCF.

As compared to the average carbon stock $(285.0 \text{ t ha}^{-1})$ of the tropical forests in the world, the Terai regions of Nepal consist of a huge amount of organic carbon (479.29 t ha⁻¹) (Jina et al., 2009). But it is lower when compared with the CFs of Nepal (Charmakar et al., 2021). For the accurate estimation of forest biomass and CS, it requires a precise calculation of both AGB and BGB (Gautam and Mandal, 2016). There was considerable variation in biomass and carbon stock at different blocks. There is variation in the size of the trees in different blocks and CS was found different. The BCCF is a growing forest with small-diameter trees. Thus, the biomass and carbon content in this forest is found quite low as compared to other studies in Shorea robusta forest and tropical forest. There is a need to carry studies on a site-specific basis representing different types of forest and conduct growth modeling to ensure the future amount of carbon.

CONCLUSION

This study has assessed the biomass and CS in the BCCF which was estimated to be 184.4 t ha⁻¹ and 99.02 t ha⁻¹ respectively. A total of 143.51 t ha⁻¹ carbon was found in block 5, which was the highest

among all blocks. The lowest (49.78 t ha⁻¹) carbon stock was found in block 6. The difference was found due to the presence of trees of higher diameters, with a higher number of trees per ha. The biomass and CS in the BCCF were found distributed in the order of tree, root, and sapling. The forest is concluded to be a growing forest having a relatively small diameter and height; it is suggested to properly manage the forest with different management operations as it has a high potential to sequester a large amount of carbon in the future. This study will help to design the blockwise different potential management inventions which can be done to enhance the quality and productivity of the forest in the future.

REFERENCES

- Adame, P., Cañellas, I., Moreno-Fernández, D., Packalen, T., Hernández, L., & Alberdi, I. (2020). Analyzing the Joint Effect of Forest Management and Wildfires on Living Biomass and Carbon Stocks in Spanish Forests. *Forests*, 11 (11), 1219.
- Alexandrov, G. A. (2007). Carbon stock growth in a forest stand: the power of age. *Carbon Balance and Management*, 2 (1), 1-5.
- ANSAB. (2010). Forest Carbon Stock Measurement: Guidelines for measuring carbon stocks in community-managed forests.
- Avitabile, V., Herold, M., Heuvelink, G.B.M., Lewis, S.L., Phillips, O.L., et al., (2016). An integrated pan-tropical biomass map using multiple reference datasets. *Global Change Biology*, 22, 1406–1420.
- Banik, B., Deb, D., Deb, S., & Datta, B. K. (2018). Assessment of biomass and carbon stock in sal (Shorea robusta Gaertn.) forests under two management regimes in Tripura, Northeast India. *Journal of forest and environmental science*, 34 (3), 209-223.
- Banskota, K., Karky, B., & Skutsch, M. (2007). Reducing carbon emissions through community-managed forests in the Himalaya. International Centre for Integrated Mountain Development (ICIMOD).
- Baral, S. K., Malla, R., & Ranabhat, S. (2009). Above-ground carbon stock assessment in different forest types of Nepal. *Banko Janakari*, 19 (2), 10-14.

- Behera, S. K., Sahu, N., Mishra, A. K., Bargali, S. S., Behera, M. D., & Tuli, R. (2017). Aboveground biomass and carbon stock assessment in Indian tropical deciduous forest and relationship with stand structural attributes. *Ecological Engineering*, 99, 513-524.
- Bhusal, M. (2011). An Analytical Study of Carbon Stock in Shores robusta Forest of Two Different Ecological Zones of Nepal (case study from Gorkha and Chitwan district). *Tribhuvan University, Pokhara*.
- Bohara, B., Miya, M. S., Timilsina, S., Gautam, D., & Regmi, S. (2021). Biomass and Carbon Stock Variation along slopes in Tropical Forest of Nepal: A case of Depard Community Forest, Makwanpur, Nepal. *Journal of Multidisciplinary Applied Natural Science*, 1(2), 89-99.
- Brown, S. L., Schroeder, P., & Kern, J. S. (1999). Spatial distribution of biomass in forests of the eastern USA. *Forest Ecology and Management*, 123 (1), 81-90.
- Canadell, J. G., & Raupach, M. R. (2008). Managing forests for climate change mitigation. *Science*, 320 (5882), 1456-1457.
- Cao, J., Wang, X., Tian, Y., Wen, Z., & Zha, T. (2012). Pattern of carbon allocation across three different stages of stand development of a Chinese pine (Pinus tabulaeformis) forest. *Ecological Research*, 27 (5), 883-892.
- CBS. (2012). National population and housing census 2011. *National Report*.
- Charmakar, S., Oli, B. N., Joshi, N. R., Maraseni, T. N., & Atreya, K. (2021). Forest carbon storage and species richness in FSC certified and non-certified community forests in Nepal. *Small-scale Forestry*, 20 (2), 199-219.
- Chave, J., Andalo, C., Brown, S., Cairns, M. A., Chambers, J. Q., Eamus, D., Fölster, H., Fromard, F., Higuchi, N., Kira, T., Lescure, J.P., Nelson, B.W., Ogawa, H., Puig, H., Riéra, B., & Yamakura, T. (2005). Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145 (1), 87-99.
- Dar, J. A., Rather, M. Y., Subashree, K., Sundarapandian, S., & Khan, M. L. (2017).
 Distribution patterns of tree, understorey, and detritus biomass in coniferous and broad-

leaved forests of Western Himalaya, India. *Journal of Sustainable Forestry*, 36 (8), 787-805.

- Dixon, R. K., Solomon, A. M., Brown, S., Houghton, R. A., Trexier, M. C., & Wisniewski, J. (1994). Carbon pools and flux of global forest ecosystems. *Science*, 263 (5144), 185-190.
- Eggleston, S., Buendia, L., Miwa, K., Ngara, T., & Tanabe, K. (2006). IPCC guidelines for national greenhouse gas inventories.
- Eriksson, E., & Berg, S. (2007). Implications of environmental quality objectives on the potential of forestry to reduce net CO2 emissions–a case study in central Sweden. *Forestry*, 80 (2), 99-111.
- Fahey, T. J., Woodbury, P. B., Battles, J. J., Goodale, C. L., Hamburg, S. P., Ollinger, S. V., & Woodall, C. W. (2010). Forest carbon storage: ecology, management, and policy. *Frontiers in Ecology and the Environment*, 8 (5), 245-252.
- FAO. (2011). FAO Glossary of Terms,. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO).
- Forsell, N., Korosuo, A., Fedeirici, S., Gusti, M., Rincón-Cristóbal, J. J., Rüter, S., Sánchez-Jiménez, B., Dore, C., Brajterman, O., & Gardiner, J. (2018). Guidance on developing and reporting the Forest Reference Levels in accordance with Regulation (EU) 2018/841.
- Gautam, K. H., & Devoe, N. N. (2006). Ecological and anthropogenic niches of sal (Shorea robusta Gaertn. f.) forest and prospects for multiple-product forest management - A review. *Forestry*, 79(1), 81–101,
- Gautam, T. P., & Mandal, T. N. (2016). Effect of disturbance on biomass, production and carbon dynamics in moist tropical forest of eastern Nepal. *Forest Ecosystems*, 3 (1), 1-10.
- Ghimire, P. (2017). *Plant Species Diversity and Tree Carbon Stock in Danphe community Forest, Kuirepani, Dang, Nepal* (Doctoral dissertation, Central Department of Botany Tribhuvan University Kirtipur, Kathmandu, Nepal).
- Gibbs, H. K., Brown, S., Niles, J. O., & Foley, J. A. (2007). Monitoring and estimating tropical

forest carbon stocks: making REDD a reality. *Environmental research letters*, 2 (4), 045023.

- Harishma, K. M., Sandeep, S., & Sreekumar, V. B. (2020). Biomass and carbon stocks in mangrove ecosystems of Kerala, southwest coast of India. *Ecological Processes*, 9 (1), 1-9.
- Houghton, R. A. (2005). Aboveground forest biomass and the global carbon balance. *Global change biology*, 11 (6), 945-958.
- IPCC. (2007). Summary for Policymakers. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. New York Cambridge University Press, Geneva. https://doi.org/10.1038/446727a
- Jackson, J. K. (1994). Manual of afforestation in Nepal, Volume 2. p.214. <u>http://www.docdeveloppement-durable.org/file/Arbres-Boisde-Rapport-</u> Reforestation/FICHES_ARBRES/bambou/Ja

cksonbamboo.pdf accessed on 20 May 2021.

- Jina, B. S., Sah, P., Bhatt, M. D., & Rawat, Y. S. (2008). Estimating carbon sequestration rates and total carbon stockpile in degraded and non-degraded sites of Oak and Pine forest of Kumaun Central Himalaya. *Ecoprint: An International Journal of Ecology*, 15, 75-81.
- Karki, S., Joshi, N. R., Udas, E., Adhikari, M. D., Sherpa, S., Kotru, R., Karky, B. S., Chhetri, N., & Ning, W. (2016). Assessment of forest carbon stock and carbon sequestration rates at the ICIMOD Knowledge Park at Godavari. *ICIMOD Working Paper*, (2016/6).
- Kaul, M., Mohren, G. M. J., & Dadhwal, V. K. (2010). Carbon storage and sequestration potential of selected tree species in India. *Mitigation and Adaptation Strategies for Global Change*, 15 (5), 489-510.
- Ketterings, Q. M., Coe, R., van Noordwijk, M., & Palm, C. A. (2001). Reducing uncertainty in the use of allometric biomass equations for predicting above-ground tree biomass in mixed secondary forests. *Forest Ecology and management*, 146 (1-3), 199-209.
- Kirschbaum, M. U. (2000). What contribution can tree plantations make towards meeting Australia's commitments under the Kyoto

Protocol?. *Environmental Science & Policy*, 3 (2-3), 83-90.

- Komolafe, E. T., Chukwuka, K. S., Obiakara, M. C., & Osonubi, O. (2020). Carbon stock and sequestration potential of Ibodi monkey forest in Atakumosa, Osun state, Nigeria. *Trees, Forests and People*, 2, 100031.
- Lewis, S. L., Sonké, B., Sunderland, T., Begne, S. K., Lopez-Gonzalez, G., Van Der Heijden, G. M. et al. (2013). Above-ground biomass and structure of 260 African tropical forests. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625), 20120295.
- Mandal, R. A., Dutta, I. C., Jha, P. K., & Karmacharya, S. (2013). Relationship between carbon stock and plant biodiversity in collaborative forests in Terai, Nepal. *International Scholarly Research Notices*, 2013.
- Mitchard, E.T.A., Feldpausch, T.R., Brienen, R.J.W., Lopez-Gonzalez, G., Monteagudo, A., et al., (2014). Markedly divergent estimates of Amazon forest carbon density from ground plots and satellites. *Global Ecology and Biogeography*, 23, 935–946.
- Mokany, K., Raison, R. J., & Prokushkin, A. S. (2006). Critical analysis of root: shoot ratios in terrestrial biomes. *Global change biology*, 12 (1), 84-96.
- Nepal, S. (2006). A Comparative study on Carbon Sequestration from two forest types in Community Forestry System (A case study from coniferous and broad leaved forests in Palpa District) B. Sc thesis submitted to Tribhuwan University, Institute of Forestry.
- Ngo, K. M., Turner, B. L., Muller-Landau, H. C., Davies, S. J., Larjavaara, M., bin Nik Hassan, N. F., & Lum, S. (2013). Carbon stocks in primary and secondary tropical forests in Singapore. *Forest Ecology and Management*, 296, 81-89.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Stephen Sitch, S., & Hayes, D. (2011). A large and persistent carbon sink in the world's forests. *Science*, 333(6045), 988-993.

- Pan, Y., Birdsey, R. A., Phillips, O. L., & Jackson, R. B. (2013). The structure, distribution, and biomass of the world's forests. *Annual Review of Ecology, Evolution, and Systematics*, 44, 593-622.
- Panday, D., Ojha, R. B., Chalise, D., Das, S., & Twanabasu, B. (2019). Spatial variability of soil properties under different land use in the Dang district of Nepal. *Cogent Food & Agriculture*, 5 (1), 1600460.
- Pandey, H. P., & Bhusal, M. (2016). A comparative study on carbon stock in Sal (Shorea robusta) forest in two different ecological regions of Nepal. *Banko Janakari*, 26 (1), 24-31.
- Pandey, S. S., Maraseni, T. N., & Cockfield, G. (2014). Carbon stock dynamics in different vegetation dominated community forests under REDD+: a case from Nepal. *Forest Ecology and Management*, 327, 40-47.
- Panzou, G. J. L., Fayolle, A., Feldpausch, T. R., Ligot, G., Doucet, J. L., Forni, E., Zombo, I., Mazengue, M., Loumeto, J. J., & Gourlet-Fleury, S. (2018). What controls local-scale aboveground biomass variation in central Africa? Testing structural, composition and architectural attributes. *Forest Ecology and Management*, 429, 570-578.
- Phillips, O. L., Malhi, Y., Higuchi, N., Laurance, W. F., Núnez, P. V., Vásquez, R. M., Laurance, S. G., Ferreira, L. V., Stern, M., Brown, S., & Grace, J. (1998). Changes in the carbon balance of tropical forests: evidence from long-term plots. *Science*, 282 (5388), 439-442.
- Raha, D., Dar, J. A., Pandey, P. K., Lone, P. A., Verma, S., Khare, P. K., & Khan, M. L. (2020). Variation in tree biomass and carbon stocks in three tropical dry deciduous forest types of Madhya Pradesh, India. *Carbon Management*, 11 (2), 109-120.
- Reichstein, M., Bahn, M., Ciais, P., Frank, D., Mahecha, M. D., Seneviratne, S. I., Zscheischler, J., Beer, C., Buchmann, N., Frank, D. C., Papale, D., Rammig, A., Smith, P., Thonicke, K., Velde, M., Vicca, S., Walz, A., & Wattenbach, M. (2013). Climate extremes and the carbon cycle. *Nature*, 500 (7462), 287-295.
- Saatchi, S.S., Harris, N.L., Brown, S., Lefsky, M., Mitchard, E.T., Salas, W., et al., (2011).

Benchmark map of forest carbon stocks in tropical regions across three continents. *PNAS* 108 (24), 9899–9904.

- Shrestha, B. (2008). An Analytical Study of Carbon Sequestration in Three Different Forest Types of Mid-Hills of Nepal. M.Sc. *Thesis*, Tribhuvan University, Institute of Forestry, Pokhara, Nepal.
- Skutsch, M., & Van Laake, P. E. (2008). Redd as multi-level governance in-the-making. *Energy & environment*, 19 (6), 831-844.
- Subedi, B. P., Pandey, S. S., Pandey, A., Rana, E. B., Bhattarai, S., Banskota, T. R., ... & Tamrakar, R. (2010). Forest carbon stock measurement: guidelines for measuring carbon stocks in community-managed forests. *Kathmandu: ANSAB, FECOFUN, ICIMOD.*
- Suryawanshi, M. N., Patel, A. R., Kale, T. S., & Patil, P. R. (2014). Carbon sequestration potential of tree species in the environment of North Maharashtra University Campus, Jalgaon (MS) India. *Bioscience Discovery*, 5 (2), 175-179.
- Tamrakar, P. R. (2000). Biomass and volume tables with species description for community forest management. His Majesty's Government of Nepal, Ministry of Forests and Soil Conservation. Natural Resource Management Sector Assistance Programme (NARMSAP), Tree Improvement and Silviculture Component.
- Tripathi, S., Thapa, C. B., & Sharma, A. (2017). Biomass Carbon Content in Schima-Castanopsis Forest of Midhills of Nepal: A Case Study from Jaisikuna Community Forest, Kaski. *Himalayan Biodiversity*, 5 (1), 1-8.
- UNFCC Kyoto Protocol Reference Manual. (2008). United Nations Framework Convention on Climate Change. <u>https://unfccc.int/sites/default/files/08_unfcc</u> <u>c_kp_ref_manual.pdf</u> accessed on 4 June 2021.
- UNFCCC. (2015). Conference of the Parties Twenty-first session, *Adoption of the Paris Agreement*. https://doi.org/10.1007/BF0232712**v**
- Upadhyay, T. P., Sankhayan, P. L., & Solberg, B. (2005). A review of carbon sequestration

dynamics in the Himalayan region as a function of land-use change and forest/soil degradation with special reference to Nepal. *Agriculture, ecosystems & environment*, 105 (3), 449-465.

- Vachnadze, G., Tiginashvili, Z., Tsereteli, G., Aptsiauri, B., & Basilidze, L. (2018). Carbon stock sequestered in the phytocenosis of oak forests in Georgia. *Annals of Agrarian Science*, 16 (4), 476-480.
- Webb, E. L., & Sah R. N. (2003). Structure and diversity of natural and managed sal (Shorea robusta Gaertn.f) forest in the Terai of Nepal. *Forest Ecology and Management*, 176 (1-3), 337–353.